

ON CONTINUOUS FIELDS OF JB-ALGEBRAS

ALEXANDER A. KATZ

Dedicated to the memory of Professor George Bachman, Polytechnic University, Brooklyn, NY, USA.

ABSTRACT. We introduce and study continuous fields of JB-algebras (which are real non-associate analogues of C^* -algebras). In particular, we show that for the universal enveloping C^* -algebra $C_u^*(B)$ for the JB-algebra B defined by a continuous field of JB-algebras $A_t, t \in T$, on a locally compact space T there exists a decomposition of $C_u^*(B)$ into a continuous field of C^* -algebras $C_u^*(A_t), t \in T$, on the same space T , composed entirely of the universal enveloping C^* -algebras of the corresponding JB-algebras from the aforementioned decomposition of the algebra B .

1. INTRODUCTION AND PRELIMINARIES

Banach associative regular $*$ -algebras over \mathbb{C} , so called *C^* -algebras*, were first introduced by Gelfand and Naimark in the paper [5]. Since then these algebras were studied extensively by various authors. This theory now is a big subdomain of the of Functional Analysis as a subject which found applications in almost all branches of Modern Mathematics and Physics. For the basics of the theory of C^* -algebras, see for example Pedersen's monograph [9]. The basic theory of real associative analogues of C^* -algebras, so called *real C^* -algebras*, is presented in Li's monograph [8].

In order to obtain a topological non-commutative version of Gelfand's characterization of commutative C^* -algebras, Dixmier and Douady in [4] introduced a notion of continuous fields of Banach spaces and C^* -algebras, which found important applications in classification of C^* -algebras (see [3]) and Theoretical Physics (see [7]). According to them, **a continuous field of C^* -algebras**

$$(\mathfrak{B}, \{\mathfrak{A}_t, \varphi_t\}_{t \in T}),$$

over a locally compact Hausdorff space T consists of a C^* -algebra \mathfrak{B} , a collection of C^* -algebras

$$\{\mathfrak{A}_t\}_{t \in T},$$

and a set

$$\{\varphi_t : \mathfrak{B} \rightarrow \mathfrak{A}_t\}_{t \in T},$$

of surjective morphisms, such that:

1). The function

$$t \mapsto \|\varphi_t(x)\|,$$

Date: October 13-14, 2007.

2000 Mathematics Subject Classification. Primary 46L, 46H; Secondary 46L70, 46H05.

Key words and phrases. C^* -algebras, real C^* -algebras, JB-algebras, continuous fields of Banach spaces, universal enveloping C^* -algebra for a JB-algebra.

is in $C_0(T)$ for all $x \in \mathfrak{B}$;

2). The norm of any $x \in \mathfrak{B}$ is

$$\|x\|_{\mathfrak{B}} = \sup_{t \in T} \|\varphi_t(x)\|_{\mathfrak{A}_t};$$

3). For any $f \in C_0(T)$ and $x \in \mathfrak{B}$, there is an element

$$fx \in \mathfrak{B},$$

for which

$$\varphi_t(fx) = f(t)\varphi_t(x),$$

for all $t \in T$.

A **section** of the field is an element $\{x_t\}_{t \in T}$ of

$$\prod_{t \in T} \mathfrak{A}_t,$$

for which there is an element $x \in \mathfrak{B}$ such that

$$x_t = \varphi_t(x),$$

for all $t \in T$.

One can see that \mathfrak{B} can be identified with the space of sections of the field, seen as a C^* -algebra under pointwise scalar multiplication, addition, adjointing, and operator multiplication, by means

$$\{\varphi_t(x)\}_{t \in T} \leftrightarrow x.$$

In particular,

$$x = y,$$

iff

$$\varphi_t(x) = \varphi_t(y),$$

for all t . It is natural that algebra \mathfrak{B} is called a the **C^* -algebra of the continuous field of C^* -algebras**.

Since the beginning of the theory of complex C^* -algebras, there were numerous attempts to extend this theory to non-associative algebras which are close to associative, in particular to Jordan algebras. In fact, Alfsen, Shultz and Størmer in [2] defined so called *JB-algebras* as the real Banach–Jordan algebras satisfying for all pairs of elements x and y the inequality of fineness

$$\|x^2 + y^2\| \geq \|x\|^2,$$

and regularity condition

$$\|x^2\| = \|x\|^2.$$

If \mathfrak{A} is a C^* -algebra, or a real C^* -algebra, then the self-adjoint part \mathfrak{A}_{sa} of \mathfrak{A} is a JB-algebra under the Jordan product

$$x \circ y = \frac{(xy + yx)}{2}.$$

Closed subalgebras of \mathfrak{A}_{sa} , for some C^* -algebra or real C^* -algebra \mathfrak{A} , become relevant examples of JB-algebras, and are called *JC-algebras*.

The basic theory of JB-algebras is fully treated in monograph of Hanche-Olsen and Størmer [6]. In particular, in this monograph there is the following theorem which was for the first time presented by Alfsen, Hanche-Olsen and Shultz in the paper [1].

Theorem 1 (Alfsen, Hanche-Olsen, Shultz [1]). *For an arbitrary JB-algebra A there exists a unique up to an isometric $*$ -isomorphism a C^* -algebra $C_u^*(A)$ (the universal enveloping C^* -algebra for the JB-algebra A), and a Jordan homomorphism*

$$\psi_A : A \rightarrow C_u^*(A)_{sa},$$

from A to the self-adjoint part of $C_u^(A)$, such that:*

- 1). $\psi_A(A)$ generates $C_u^*(A)$ as a C^* -algebra;
- 2). for any pair composed of a C^* -algebra \mathfrak{A} and a Jordan homomorphism

$$\rho : A \rightarrow \mathfrak{A}_{sa},$$

from A into the self-adjoint part of \mathfrak{A} , there exists a $$ -homomorphism*

$$\hat{\rho} : C_u^*(A) \rightarrow \mathfrak{A},$$

from the C^ -algebra $C_u^*(A)$ into C^* -algebra \mathfrak{A} , such that*

$$\rho = \hat{\rho} \circ \psi_A;$$

- 3). *there exists a $*$ -antiautomorphism Φ of order 2 on the C^* -algebra $C_u^*(A)$, such that*

$$\Phi(\psi_A(a)) = \psi_A(a),$$

$\forall a \in A$.

□

Our plan is to define a *continuous field of JB-algebras*, the *JB-algebra of the continuous field of JB-algebras*, and be able in the spirit of Theorem 1 above to associate in a universal sense with each JB-algebra of the continuous field of JB-algebras a C^* -algebra of the continuous field of C^* -algebras.

2. CONTINUOUS FIELDS OF JB-ALGEBRAS

Let us first introduce a continuous field of JB-algebras.

Definition 1. *A continuous field of JB-algebras*

$$(B, \{A_t, \varphi_t\}_{t \in T}),$$

over a locally compact Hausdorff space T consists of a JB-algebra B , a collection of JB-algebras $\{A_t\}_{t \in T}$, and a set

$$\{\varphi_t : B \rightarrow A_t\}_{t \in T},$$

of surjective morphisms, such that:

- 1). *The function*

$$t \mapsto \|\varphi_t(x)\|,$$

is in $C_0(T)$ for all $x \in B$;

- 2). *The norm of any $x \in B$ is*

$$\|x\|_B = \sup_{t \in T} \|\varphi_t(x)\|_{A_t};$$

- 3). *For any $f \in C_0(T)$ and $x \in B$, there is an element*

$$fx \in B,$$

for which

$$\varphi_t(fx) = f(t)\varphi_t(x),$$

for all $t \in T$.

A **section** of the field is an element $\{x_t\}_{t \in T}$ of

$$\prod_{t \in T} A_t,$$

for which there is an element $x \in B$ such that

$$x_t = \varphi_t(x),$$

for all $t \in T$.

We identify B with the space of sections of the field, seen as a JB-algebra under pointwise scalar multiplication, addition, operator multiplication, by means

$$\{\varphi_t(x)\}_{t \in T} \leftrightarrow x.$$

In particular,

$$x = y,$$

iff

$$\varphi_t(x) = \varphi_t(y),$$

for all t . It is natural that algebra B is called a the **JB-algebra of the continuous field of JB-algebras**.

Now we will establish a few properties of the continuous field of JB-algebras. The first one is about locally uniform closedness of the continuous field of JB-algebras.

Proposition 1. *The JB-algebra B of sections of a continuous field of JB-algebras is **locally uniformly closed**, i.e. if*

$$x \in \prod_{t \in T} A_t,$$

is such that for every

$$s \in T,$$

and every $\varepsilon > 0$ there exists

$$y_s \in B,$$

and a neighborhood

$$V_s \subset T,$$

of s in which

$$\|x_t - \varphi_t(y_s)\| < \varepsilon,$$

for all

$$t \in V_s,$$

and also

$$\lim_{t \rightarrow \infty} \|x_t\| = 0,$$

then

$$x \in B.$$

Alternatively, if the function

$$t \mapsto \|x_t - z_t\|,$$

lies in $C_0(T)$ for each

$$z \in B,$$

then

$$x \in B.$$

Proof. Under conditions of the first part of the Proposition, there exists a compact set

$$K \subseteq T,$$

for which

$$\|x_t\| < \varepsilon,$$

outside of K , as well as a finite cover

$$\{V_{t_1}, \dots, V_{t_n}\},$$

of K ,

$$K \subseteq \{V_{t_1}, \dots, V_{t_n}\}.$$

Now we have to recall a notion of a partition of unity on K subordinate to this cover (see for example [10] and [7]). Let K be a Hausdorff space, and

$$\{V_\alpha\}_{\alpha \in \Lambda},$$

be a **locally finite open cover** of K , i.e. each point of K has a neighborhood that intersects only with a finite number of the sets V_α . A **partition of unity subordinate to the given cover** is a collection of positive functions

$$\{u_\alpha\}_{\alpha \in \Lambda},$$

such that each u_α is a compactly supported continuous real-valued function with

$$\sum_{\alpha \in \Lambda} u_\alpha = 1.$$

A partition of unity always exists when K is paracompact (see [10]). So, let us take a partition of unity

$$\{u_i\}_{i=1}^n,$$

on K subordinate to the aforementioned finite cover

$$\{V_{t_1}, \dots, V_{t_n}\}.$$

Let us consider the

$$y = \sum_{i=1}^n u_i y_{t_i}.$$

From Definition 1.3 it follows that

$$y \in B,$$

and satisfies the condition

$$\sup_{t \in T} \|x_t - y_t\| < \varepsilon.$$

Therefore, from Definition 1.2 and completeness of B it follows that

$$x \in B.$$

Now, given any

$$x \in \prod_{t \in T} A_t,$$

and

$$s \in T,$$

because φ_s is surjective, there exists an element

$$y_s \in B,$$

such that

$$x_s = \varphi_s(y_s).$$

The assumption of the second part of Proposition 1 then implies that the conditions in the first part of this Proposition are satisfied, such that

$$x \in B.$$

□

The following Proposition gives conditions for the existence and uniqueness of a continuous field of JB-algebras whose collection of sections contains a subset possessing some natural properties.

Proposition 2. *Let*

$$\{A_t\}_{t \in T},$$

be a family of JB-algebras indexed by a locally compact Hausdorff space T , and a subset

$$\tilde{B} \subseteq \prod_{t \in T} A_t,$$

that satisfies the following properties:

1). *The set*

$$\{x_t : x \in \tilde{B}\},$$

is dense in A_t for each $t \in T$;

2). *The function*

$$t \mapsto \|x_t\|,$$

lies in $C_0(T)$ for each $x \in \tilde{B}$;

3). *The set \tilde{B} is a Jordan algebra under pointwise operations.*

Then there exists a unique continuous field of JB-algebras

$$(B, \{A_t, \varphi_t\}_{t \in T}),$$

whose collection of sections contains \tilde{B} . Namely, B consists of all

$$x \in \prod_{t \in T} A_t,$$

for which the function

$$x \mapsto \|x_t - z_t\|,$$

lies in $C_0(T)$ for each $z \in \tilde{B}$, regarded as JB-algebra under pointwise operations, and the norm of Definition 1.2. Finally,

$$\varphi_t(x) = x_t,$$

$t \in T$, is the evaluation map.

Proof. We show first that the algebra B defined above is locally uniformly closed. With the objects x, s, ε, y_s and V as specified in Proposition 1, take $z \in \tilde{B}$ arbitrary, and define the functions

$$f_{xz} : t \mapsto \|x_t - z_t\|,$$

and

$$f_{yz} : t \mapsto \|\varphi_t(y_s) - z_t\|.$$

Using the triangle inequality for the norm in Banach space, we get that

$$|(\|x\| - \|y\|)| \leq \|x - y\|,$$

and that gives us that

$$|f_{xz}(t) - f_{yz}(t)| < \varepsilon,$$

for all $t \in V$. By assumption, the function f_{yz} is continuous, so that

$$|f_{yz}(t) - f_{yz}(s)| < \varepsilon,$$

for all t 's in some neighborhood V' of s . Combining the inequalities, we get

$$|f_{xz}(t) - f_{xz}(s)| < 3\varepsilon,$$

for all

$$t \in V \cap V'.$$

Therefore f_{xz} is continuous at s , which was arbitrary, so that $x \in B$ by the definition of B .

Now, we show uniqueness of B . Using this property one can easily see that B is a JB-algebra, and that the condition 3 in Definition 1 is satisfied. It is clear from Definition 1.1 and the definition of B in Proposition 2 that B is maximal. On the other hand, according to the second part of Proposition 1, B is minimal, so, B is unique. \square

We are ready now to present the main result of the paper.

Definition 2. An $*$ -isomorphism (resp. Jordan isomorphism) of continuous fields of C^* -algebras (resp. JB-algebras) over the same base Hausdorff locally compact topological space T is the isometric $*$ -isomorphism (Jordan isometric isomorphism) of the C^* -algebras (resp. JB-algebras) of the continuous fields via a map respecting the fibers.

Theorem 2. For an arbitrary continuous field of JB-algebras

$$(B, \{A_t, \varphi_t\}_{t \in T}),$$

over a locally compact Hausdorff space T , there exists a unique up to an $*$ -isomorphism a continuous field of C^* -algebras

$$(C_u^*(B), \{C_u^*(A_t), \widehat{\varphi}_t\}_{t \in T}),$$

(the universal enveloping continuous field of C^* -algebras for the continuous field of JB-algebras

$$(B, \{A_t, \varphi_t\}_{t \in T}))$$

over the same base space T), and a Jordan homomorphism

$$\psi_B : B \rightarrow C_u^*(B)_{sa},$$

from B to the self-adjoint part of $C_u^*(B)$, as well as a family of Jordan homomorphisms

$$\psi_{A_t} : A_t \rightarrow C_u^*(A_t)_{sa},$$

$t \in T$, from A_t to the self-adjoint part of $C_u^*(A_t)$, for each $t \in T$, such that:

- 1). $\psi_B(B)$ generates $C_u^*(B)$ as a C^* -algebra, and each $\psi_{A_t}(B)$ generates each $C_u^*(A_t)$ as a C^* -algebra for each $t \in T$;
- 2). for any pair composed of a continuous field of C^* -algebras $(\mathfrak{B}, \{\mathfrak{A}_t, \widetilde{\varphi}_t\}_{t \in T})$ and a family of Jordan homomorphisms

$$\rho : B \rightarrow \mathfrak{B}_{sa},$$

from B into the self-adjoint part of \mathfrak{B} , and

$$\rho_t : A_t \rightarrow (\mathfrak{A}_t)_{sa},$$

for each $t \in T$, from A_t into the self-adjoint part of \mathfrak{A}_t , there exist a $*$ -homomorphism

$$\widehat{\rho} : C_u^*(B) \rightarrow \mathfrak{B},$$

from the C^* -algebra $C_u^*(B)$ into C^* -algebra \mathfrak{B} , and a family of $*$ -homomorphisms

$$\widehat{\rho}_t : C_u^*(A_t) \rightarrow \mathfrak{A}_t,$$

for each $t \in T$, from the C^* -algebra $C_u^*(A_t)$ into C^* -algebra \mathfrak{A}_t such that

$$\rho = \widehat{\rho} \circ \psi_B,$$

and

$$\rho_t = \widehat{\rho}_t \circ \psi_{A_t},$$

for each $t \in T$;

3). there exists a $*$ -antiautomorphism Φ of order 2 on the C^* -algebra $C_u^*(B)$, such that

$$\Phi(\psi_B(x)) = \psi_B(x),$$

$\forall x \in B$, as well as there exists a family of $*$ -antiautomorphism Φ_t of order 2 on the C^* -algebra $C_u^*(A_t)$, for each $t \in T$, such that

$$\Phi(\psi_{A_t}(x_t)) = \psi_{A_t}(x_t),$$

$\forall x_t \in A_t$, and every $t \in T$.

Proof. Let $(B, \{A_t, \varphi_t\}_{t \in T})$ be a given continuous field of JB-algebras. Let $C_u^*(B)$ be the universal enveloping C^* -algebra for the JB-algebra of the continuous field, and the family of C^* -algebras $C_u^*(A_t)$ for each $t \in T$ be the universal enveloping C^* -algebra for the JB-algebra A_t , $t \in T$. Let

$$\psi_B : B \rightarrow C_u^*(B)_{sa},$$

and for each $t \in T$,

$$\varphi_t : B \rightarrow A_t,$$

and

$$\psi_{A_t} : A_t \rightarrow C_u^*(A_t)_{sa}.$$

From Theorem 1 it follows that $\psi_B(B)$ is dense in $C_u^*(B)_{sa}$, and $\psi_{A_t}(A_t)$ is dense in $C_u^*(A_t)_{sa}$ for each $t \in T$. So, without a loss of generality we can assume that for each

$$x, y \in C_u^*(B)_{sa}$$

there exist $a_n, b_n \in B$, such that

$$x = \lim_{n \rightarrow \infty} \psi_B(a_n),$$

and

$$y = \lim_{n \rightarrow \infty} \psi_B(b_n),$$

where the limit is taken in the norm of $C_u^*(B)$, as well as

$$x_t = \widehat{\varphi}_t(x) = \lim_{n \rightarrow \infty} \psi_{A_t}(\varphi_t(a_n)),$$

and

$$y_t = \widehat{\varphi}_t(y) = \lim_{n \rightarrow \infty} \psi_{A_t}(\varphi_t(b_n)),$$

$t \in T$, where the limit is taken in the norm of $C_u^*(A_t)$. For each $t \in T$, we will define

$$\widehat{\varphi}_t : C_u^*(B) \rightarrow C_u^*(A_t),$$

the following way:

$$\widehat{\varphi}_t(x + iy) = x_t + iy_t.$$

Because $C_u^*(A_t)_{sa}$ is norm closed for each $t \in T$ (see [9]), the last identity is well defined. Moreover, from the fact that φ_t is surjective for each $t \in T$ it follows that $\widehat{\varphi}_t$ is surjective for each $t \in T$ as well. Thus,

$$(C_u^*(B), \{C_u^*(A_t), \widehat{\varphi}_t\}_{t \in T}),$$

is in fact a continuous field of C^* -algebras. The rest of the Theorem is obtained as a corollary by application of Theorem 1 in fibers, and Propositions 1 and 2. \square

REFERENCES

- [1] **Alfsen, E.M.; Hanche-Olsen, H.; Shultz, F.W.**, *State spaces of C^* -algebras*. (English) Acta Math. Vol. 144 (1980), No. 3-4, pp. 267-305.
- [2] **Alfsen, E.M.; Shultz, F.W.; Størmer, E.**, *A Gelfand-Naimark theorem for Jordan algebras*. (English) Advances in Math. Vol. 28 (1978), No. 1, pp. 11-56.
- [3] **Dixmier, J.**, *C^* -algebras*. (English, Translated from French) North-Holland Mathematical Library, Vol. 15. North-Holland Publishing Co., Amsterdam-New York-Oxford, 492 pp., (1977).
- [4] **Dixmier, J.; Douady, A.**, *Champs Continus d'Espaces Hilbertiens et de C^* -Algebres*. (French) Bull. Soc. math. France Vol. 91 (1963), pp. 227-284.
- [5] **Gelfand, I.M.; Naimark, M.A.**, *On the embedding of normed rings into the ring of operators in Hilbert space*. (English. Russian summary) Rec. Math. [Mat. Sbornik] N.S. Vol. 12(54) (1943), pp. 197-213.
- [6] **Hanche-Olsen, H.; Størmer, E.**, *Jordan operator algebras*. (English), Monographs and Studies in Mathematics, Vol. 21. Boston - London - Melbourne: Pitman Advanced Publishing Program. VIII, 183 pp., (1984).
- [7] **Landsman, N.P.**, *Mathematical topics between classical and quantum mechanics*. (English), Springer Monographs in Mathematics. Springer-Verlag, New York, 529 pp., (1998).
- [8] **Li, B.**, *Real operator algebras*. (English) World Scientific Publishing Co., Inc., River Edge, NJ, 241 pp., (2003).
- [9] **Pedersen, G.K.**, *C^* -algebras and their automorphism groups*. (English), London Mathematical Society Monographs. Vol. 14. London - New York - San Francisco: Academic Press., 416 pp., (1979).
- [10] **Pedersen, G.K.**, *Analysis now*. (English) Graduate Texts in Mathematics, Vol. 118. Springer-Verlag, New York, 277 pp., (1989).

DR. ALEXANDER A. KATZ, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S COLLEGE OF LIBERAL ARTS AND SCIENCES, ST. JOHN'S UNIVERSITY, 300 HOWARD AVENUE, DASILVA ACADEMIC CENTER 314, STATEN ISLAND, NY 10301, USA

E-mail address: `katza@stjohns.edu`